The field of the electromagnetic knot and the Sparling—Tod metrics

© V.N. Trishin

Bauman Moscow State Technical University, Moscow, 105005, Russia

The article considers explicit anti-selfdual solutions of complex Einstein equations constructed from null Maxwell fields. It is shown that the solution of source-free Maxwell equations describing electromagnetic unit corresponds to well-known Sparling—Tod metrics.

REFERENCES

[1] Tod K.P. Self-dual Kerr—Schild metrics and null Maxwell fields. *Journal of Mathematical Physics*, 1982, vol. 23, pp. 1147–1148.

[2] Penrose R., Rindler W. *Spinors and space-time. Vol. 2. Spinor and Twistor Methods in Space-Time Geometry.* Cambridge etc., Cambridge University Press, 1986. IX, 501 p., ISBN 0 521 25267 9.

[3] Dunajski M. *Solitons, instantons, and twistors*. Oxford, OUP, 2010, 359 p.

[4] Plebanski J. Some solutions of complex Einstein equations. *Journal of Mathematical Physics*, 1975, vol. 16, pp. 2395–2402.

[5] Robinson I. Null electromagnetic fields. *Journal of Mathematical Physics*, 1961, vol. 2, pp. 290–291.

[6] Nagatomo K. On a construction of null electromagnetic fields. Osaka Journal of Mathematics, 1983, vol. 20, pp. 285–301.

[7] Dalhuisen J. W., Bouwmeester D. Twistors and electromagnetic knots. *Journal of Physics*, 2012, vol. A45, pp. 135201–135209.

[8] Sparling G.A.J., Tod K.P. An example of an H-space. *Journal of Mathematical Physics*, 1981, vol. 22, pp. 331–332.

Trishin V.N. (b. 1977) graduated from the Physics Department of Lomonosov Moscow State University. Ph.D, Assoc. Professor of the Computational Mathematics and Mathematical Physics Department of Bauman Moscow State Technical University. Author of 7 scientific papers in the field of mathematical methods in the general theory of relativity.
e-mail: trishinvn@bmstu.ru